



## Mouse obestatin ELISA kit-Quantitative competitive

<b>Cat. No.:</b>	0126WXX-1188
<b>Assay Type:</b>	Quantitative competitive ELISA
<b>Target Species:</b>	Mouse
<b>Assay Target:</b>	Obestatin
<b>Size:</b>	96T

This product is for research use only and is not intended for diagnostic use.

### Product Overview

<b>Description</b>	Mouse obestatin ELISA kit-Quantitative competitive is an ELISA-based <i>in vitro</i> research tool designed specifically for the quantitative detection of obestatin in mice samples with a range of 61.7-5000 pg/mL and a minimum detectable dose (sensitivity) of 27.5 pg/mL. The kit is highly sensitive and easy to use.
<b>Assay Principle</b>	The ELISA analytical biochemical technique is based on obestatin antibody-obestatin antigen interactions (immunosorbency) and an HRP colorimetric detection system to detect obestatin antigen targets in samples.
<b>Background</b>	Obestatin is a hormone secreted by cells in the stomach and small intestine that may regulate both hunger and satiety signals through a complex balancing system. Studying how obestatin interacts with other hormones, such as insulin and leptin, can help us gain a more comprehensive understanding of energy metabolism and analyze its role in obesity.
<b>Synonyms</b>	OB
<b>Formula Weight</b>	18,641 Da
<b>Applications</b>	Mouse obestatin ELISA kit-Quantitative competitive is used to quantify obestatin in serum, plasma, tissue homogenates, and cell lysates, cell culture supernatants, and other biological fluids of mouse, providing data to support research in a wide range of areas, including metabolic, endocrinology, obesity, etc.
<b>Research Area</b>	Metabolic; Endocrinology; Obesity

### Specification



<b>Sample Type</b>	Serum; Plasma; Tissue homogenates; Cell lysates; Cell culture supernatants; Other biological fluids
<b>Detection Range</b>	61.7-5000 pg/mL
<b>Sensitivity</b>	27.5 pg/mL
<b>Cross-reactivity</b>	No significant cross-reactivity or interference was observed.
<b>Stability</b>	12 months
<b>Storage</b>	Store at -20°C (antibodies, standard, streptavidin-HRP), 4°C (TMB) and room temperature (96-well plate).